

UNIFIED CONSTRUCTIONS OF THE REGULAR HEPTAGON AND TRISKAIDECAGON

Ruhland H., MSc, Independent Researcher, ✉ helmut.ruhland50@web.de

Abstract

Constructions of regular heptagon and triskaidecagon by trisection of an angle are well known. An elegant construction of the heptagon by S. Adlaj shows a 3-fold symmetry related to a Galois group. Based on the latter construction, in this article one more for the heptagon and two more for the triskaidecagon are presented, all using angle trisection.

Keywords: *geometric construction, trisection, heptagon, triskaidecagon, tridecagon.*

Citation: H. Ruhland, "Unified Constructions of the Regular Heptagon and Triskaidecagon", *Computer tools in education*, no. 1, pp. 27–32, 2025 (in Russian); doi:10.32603/2071-2340-2025-1-27-32

1. INTRODUCTION

In [1] A. Gleason gave constructions of a regular heptagon and triskaidecagon using only square roots and the trisection of angles. Later S. Adlaj [2] presented a very elegant construction of the heptagon, that shows the action of a cyclic subgroup of order 3 in the related Galois group on the 3 constructed vertices.

In this article I present 3 new unified constructions of the 2 regular polygons by trisection, unified means here *all* based on S. Adlaj's geometric construction. They all use angle trisection.

In section 2 I repeat the construction in [2] and show one more for the heptagon. In section 3 2 constructions for the trikaidecagon are shown.

2. CONSTRUCTIONS OF THE HEPTAGON

2.1. The first construction, type I: S. Adlaj's from [2]

Let $\varepsilon_k = e^{2\pi i/3k} = ((-1 + \sqrt{3}i)/2)^k$, $k = 0, 1, 2$ a third root of unity. Define the 3 third roots ζ_k , $k = 0, 1, 2$:

$$\zeta_k = \varepsilon_k \sqrt[3]{\zeta}, \quad \zeta = \frac{1 - 3\sqrt{3}i}{2\sqrt{7}}. \quad (1)$$

Determining the third root of ζ is a equivalent to a trisection because $|\zeta| = 1$. The angle to trisect is $\theta = -\arctan(3\sqrt{3}) \approx -79.1066^\circ$

Take as radii of the 2 grey, concentric circles in figure 1:

$$R_1, R_2 = \sqrt{(7 \pm \sqrt{21})/18}. \quad (2)$$

Form three vertices of the heptagon, the three red (—), green (---), blue (---) parallelograms in figure 1 realize the following complex additions:

$$V_0 = R_1 \varepsilon_0 + R_2 \zeta_1, \quad V_1 = R_1 \varepsilon_1 + R_2 \zeta_0, \quad V_2 = R_1 \varepsilon_2 + R_2 \zeta_2. \quad (3)$$




Figure 1. A scan of the heptagon construction in [2] with the denomination of vertices added by the author of this article. The vertices ε_k, ζ_k do not have unit distance to 0, instead they should have been denominated $\varepsilon_k R_1, \zeta_k R_2$

The 3 constructed vertices V_0, V_1, V_2 represent the quadratic residues $PR_2 = \{1, 2, 4\}$ modulo 7. The triangle built by these 3 vertices has 3 different side lengths and so no symmetry. The cyclic permutation $C_3 : \zeta_k \rightarrow \zeta_{k+1}$ of order 3, subscripts modulo 3, acts on these 3 vertices.

2.2. The second construction, Type II

Now the angle to trisect is $\theta = 0 = 0^\circ$. Only a square root is necessary to get $\theta/3 = 0^\circ, 120^\circ, 240^\circ$.

Set $R_1 = 1$ and R_2 equals one of the six real roots $r_k, k = 0, \dots, 5$ of the palindromic, sextic polynomial $P = r^6 + 6r^5 - 6r^4 - 29r^3 - 6r^2 - 6r + 1$. Because P is palindromic, with every root r , the inverse is a root too. The 6 roots can be constructed by square roots and solving a cubic, because with $Q = (s^3 + 6s^2 - 9s - 41)$ and $s = r + 1/r$, $P = Q(s)r^3$.

These 6 roots can also be expressed by the ζ_k gotten by the trisection 1:

$$s_k = -2 + \sqrt{7}(\zeta_k + \bar{\zeta}_k), \quad (4)$$

$$r_k, r_{k+3} = \frac{s_k \pm \sqrt{s_k^2 - 4}}{2}, \quad s_k \text{ a root of } Q, \quad k = 0, 1, 2. \quad (5)$$

¹There is no typing error here, in V_0, V_1 the ε and ζ subscripts are different

The vertices of the heptagon are given by 3. Hint: a negative R_2 in this geometric construction means: a vector in the parallelogram has to be taken negative.

In contrary to the previous construction, the triangle built by the 3 constructed vertices V_0, V_1, V_2 is now an *isosceles* triangle with a reflection symmetry.

$C_3 : \zeta_k \rightarrow \zeta_{k+1}$ acts now on the 6 radii r_k and so on the 6 isosceles triangles. Up to scaling and rotation are only 3 different triangles.

2.3. The third construction, but nothing new, is of type II too

Now the angle to trisect is $\theta = \pi = 180^\circ$. Only a square root is necessary to get $\theta/3 = 60^\circ, 180^\circ, 300^\circ$.

Set $R_1 = 1$ and R_2 equals the *negative* of one of the six real roots r_k of the second construction in 2.2.

Because a negative radius R_2 is equivalent to a positive radius and a rotation by π of the triangle inscribed in this circle: this construction is equivalent to the previous subsection 2.2.

3. CONSTRUCTIONS OF THE TRISKAIDECAGON

3.1. The first construction, type I

Let $\varepsilon_k = e^{2\pi i/3k} = ((-1 + \sqrt{3}i)/2)^k$, $k = 0, 1, 2$ a third root of unity. Define the 3 third roots ζ_k , $k = 0, 1, 2$:

$$\zeta_k^\pm = \varepsilon_k^3 \sqrt[3]{\zeta^\pm}, \quad \zeta^\pm = \frac{\sqrt{26 \pm 5\sqrt{13}} - \sqrt{26 \mp 5\sqrt{13}}i}{2\sqrt{13}}. \quad (6)$$

Determining the third root of ζ^\pm is a equivalent to a trisection because $|\zeta^\pm| = 1$. The 2 corresponding 2 angles to trisect are:

$$\theta^\pm = -\arctan\left(\frac{26 \mp 5\sqrt{13}}{3\sqrt{3}\sqrt{13}}\right). \quad (7)$$

Take as radii of the 2 grey, concentric circles and angle to trisect, in figure 1:

$$R_1, R_2 = \sqrt{\frac{\sqrt{13 + \sqrt{13}} \pm \sqrt{5 + \sqrt{13}}}{2\sqrt{2}}}, \quad R_1 R_2 = 1 \quad \theta^+ \approx -23.0510^\circ. \quad (8)$$

The 3 constructed vertices V_0, V_1, V_2 , see 3, represent the now quartic residues $PR_4 = \{1, 3, 9\}$ modulo 13. Using $-\theta^+/3$ the vertices represent $4PR_4 = \{4, 10, 12\}$. The triangle built by these 3 vertices has 3 different side lengths.

The cyclic permutation $C_3 : \zeta_k \rightarrow \zeta_{k+1}$ of order 3, subscripts modulo 3, acts on these 3 vertices.

To get the vertices belonging to the 2 remaining multiplicative cosets of quartic residues $2PR_4 = \{2, 5, 6\}$ and $8PR_4 = \{7, 8, 11\}$, the following 2 radii and angle to trisect have to be used:

$$R_1, R_2 = \sqrt{\frac{\sqrt{13 - \sqrt{13}} \pm \sqrt{5 - \sqrt{13}}}{2\sqrt{2}}}, \quad R_1 R_2 = 1, \quad \theta^- \approx -66.9489^\circ. \quad (9)$$

3.2. The second construction, Type II

Now the angle to trisect is $\theta = 0 = 0^\circ$. Only a square root is necessary to get $\theta/3 = 0^\circ, 120^\circ, 240^\circ$.

Set $R_1 = 1$ and R_2 equals one of the 12 real roots $r_k, k = 0, 1, \dots, 11$ of the palindromic, degree 12 polynomial $P = r^{12} + 12r^{11} - 12r^{10} - 274r^9 - 441r^8 + 441r^7 + 1275r^6 + 441r^5 - 441r^4 - 274r^3 - 12r^2 + 12r + 1$. Because P is palindromic, with every root r , the inverse is a root too. The 12 roots can be constructed by square roots and solving a cubic, because with $Q = (s^6 + 12s^5 - 18s^4 - 334s^3 - 384s^2 + 1323s + 2131)$ and $s = r + 1/r$, $P = Q(s)r^6$. The sextic Q factorizes in $\mathbb{Q}(\sqrt{13})$ as product of $R = s^3 + 3(2 + \sqrt{13})s^2 + 21(3 + \sqrt{13})/2s + (15 + 107\sqrt{13})/2$ and its $\mathbb{Q}(\sqrt{13})$ -conjugate \bar{R} .

These 12 roots can be expressed by the ζ_k^\pm gotten by the trisection 6:

$$s_k^\pm = -(2 \pm \sqrt{13}) + \sqrt{13 \pm \sqrt{13}(\zeta_k^\pm + \bar{\zeta}_k^\pm)}, \quad (10)$$

$$r_k, r_{k+3} = \frac{s_k^\pm \pm \sqrt{(s_k^\pm)^2 - 4}}{2}, \quad s_k^+ \text{ a root of } R, \quad k = 0, 1, 2, \quad (11)$$

$$r_{k+6}, r_{k+9} = \frac{s_k^\pm \pm \sqrt{(s_k^\pm)^2 - 4}}{2}, \quad s_k^- \text{ a root of } \bar{R}, \quad k = 0, 1, 2. \quad (12)$$

The conjugation on the rhs of s_k^\pm is now a \mathbb{C} -conjugation.

The vertices of the trikaidecagon are also given by 3. Hint: a negative R_2 in this geometric construction means: a vector in the parallelogram has to be taken negative.

In contrary to the previous construction, the triangle built by the 3 constructed vertices V_0, V_1, V_2 is now an *isosceles* triangle with a reflection symmetry.

$C_3 : \zeta_k \rightarrow \zeta_{k+1}$ acts now on the 12 radii r_k and so on the 12 isosceles triangles. Up to scaling and rotation are only 6 different triangles.

3.3. The third construction, but nothing new, is of type II too

Now the angle to trisect is $\theta = \pi = 180^\circ$. Only a square root is necessary to get $\theta/3 = 60^\circ, 180^\circ, 300^\circ$.

Set $R_1 = 1$ and R_2 equals the *negative* of one of the 12 real roots r_k of the second construction in 3.2.

Because a negative radius R_2 is equivalent to a positive radius and a rotation of the triangle inscribed in this circle: this construction is equivalent to the previous subsection 3.2.

4. CONSTRUCTIONS OF THE PENTAGON

4.1. The first construction, n o type I

A type I construction does not exist. Every triangle built by vertices in a regular pentagon is an isosceles triangle. A triangle with 3 different side lengths for type I does not exist.

4.2. The second construction, Type II

Now the angle to trisect is $\theta = 0 = 0^\circ$. Only a square root is necessary to get $\theta/3 = 0^\circ, 120^\circ, 240^\circ$.

Set $R_1 = 1$ and R_2 equals one of the four real roots $r_k, k = 0, \dots, 3$ of the palindromic, quartic polynomial $P = r^4 - r^3 - 9r^2 - r + 1$. Because P is palindromic, with every root r , the inverse

is a root too. The 4 roots can be constructed by square roots, because with $Q = s^2 - s - 11$ and $s = r + 1/r$, $P = Q(s)r^2$. No trisection is needed.

$$s_0, s_1 = (1 \pm 3\sqrt{5})/2 \quad \text{the 2 roots of } Q \quad (13)$$

$$r_k, r_{k+2} = \frac{s_k \pm \sqrt{s_k^2 - 4}}{2} \quad k = 0, 1 \quad (14)$$

The vertices of the pentagon are given by 3. Hint: a negative R_2 in this geometric construction means: a vector in the parallelogram has to be taken negative.

The triangle built by the 3 constructed vertices V_0, V_1, V_2 is an *isosceles* triangle with a reflection symmetry.

4.3. The third construction, but nothing new, is of type II too

Now the angle to trisect is $\theta = \pi = 180^\circ$. Only a square root is necessary to get $\theta/3 = 60^\circ, 180^\circ, 300^\circ$.

Set $R_1 = 1$ and R_2 equals the *negative* of one of the four real roots r_k of the second construction in 4.2.

Because a negative radius R_2 is equivalent to a positive radius and a rotation by π of the triangle inscribed in this circle: this construction is equivalent to the previous subsection 4.2.

5. OPEN QUESTIONS

Are the constructions of type I, type II or both also possible for other p-gons with p a prime of the form $6n + 1$? The next primes p to investigate would be $p = 19, 31, \dots$

References

1. A. Gleason, "Angle Trisection, the Heptagon, and the Triskaidecagon," *The American Mathematical Monthly*, vol. 95, no. 3, pp. 185–194, 1988; doi:10.2307/2323624
2. S. Adlaj, "The Heptagon constructed!," in *semjonadlaj.com*, 2024. [Online]. Available: <https://SemjonAdlaj.com/Excerpts/Heptagon.pdf>

Received 20-03-2025, the final version — 15-04-2025.

Helmut Ruhland, MSc, Independent Researcher, Calle 282, #1104, Santa Fe, La Habana, Cuba,
 ✉ helmut.ruhland50@web.de

Компьютерные инструменты в образовании, 2025

№ 1: 27–32

УДК: 514.172.45:514.74

<http://cte.eltech.ru>

[doi:10.32603/2071-2340-2025-1-27-32](https://doi.org/10.32603/2071-2340-2025-1-27-32)

Унифицированные построения правильных семиугольников и тринадцатиугольников

Руланд Х.¹, MSc, независимый исследователь, ✉ helmut.ruhland50@web.de

Аннотация

Построения правильных семиугольника и тринадцатиугольника путем трисекции угла хорошо известны. Элегантная конструкция семиугольника С. Ф. Адлай демонстрирует 3-кратную симметрию, связанную с группой Галуа. Основываясь на этой конструкции, в этой статье мы рассмотрим еще одну для семиугольника и еще две для тринадцатиугольника, все с использованием трисекции угла.

Ключевые слова: геометрическая конструкция, трисекция, семиугольник, тринадцатиугольник.

Цитирование: Руланд Х. Унифицированные построения правильных семиугольников и тринадцатиугольников // Компьютерные инструменты в образовании. 2025. № 1. С. 27–32. [doi:10.32603/2071-2340-2025-1-27-32](https://doi.org/10.32603/2071-2340-2025-1-27-32)

Поступила в редакцию 20.03.2025, окончательный вариант — 15.04.2025.

Хельмут Руланд, MSc, независимый исследователь, Калле 282, #1104, Санта-Фе, Гавана, Куба, ✉ helmut.ruhland50@web.de